

Have You Experienced a Stroke?

You might also experience these symptoms: stiffness, spasms, pain, and weakness. Fortunately, you have options.

Stroke is one of the most prevalent conditions, with 6.5 million people having experienced one in the US. Of those stroke sufferers, an estimated 1.5 (23%) have also experienced spasticity⁵; most often within 12 months after the stroke³.

However, everyone is different and you may experience these symptoms^{1,2} shortly following your stroke or years after.

Stiffness

Tightness or flexing of the muscles which may make it more difficult to move or perform daily activities.

Spasms

When your muscle flexes suddenly, it can cause a part of your body to jerk. This is called a spasm. Spasms happen most often in your arms, legs, hands, feet, and torso.

Pain

Stiffness and spasms can make your muscles hurt. Pain can also occur when your symptoms force you to sit or sleep in different positions.

Weakness

Weakness, fatigue and paralysis can lead to problems with balance or coordination, making it difficult to perform daily activities.

YOU'RE NOT ALONE IN THIS AND YOU HAVE OPTIONS.





A REAL PATIENT EXPERIENCE WITH SPASTICITY RELATED TO A STROKE⁴

MARY*, 72

Mary is a 72-year-old woman who suffered a stroke that affected her right upper and lower extremities. Along with weakness and fatigue, Mary developed significant spasticity, experiencing extreme tightness on her right side that resulted in poor balance. These symptoms interfered dramatically with her daily activities, like bathing, dressing and brushing her teeth. Not to mention, it affected her mobility, as she was in a wheelchair, making her transfers and ambulation difficult to achieve.

Her primary care physician referred her to a physical medicine and rehabilitation physician that prescribed oral medications including oral baclofen. When the symptoms persisted, she and her doctor agreed that **Intrathecal Baclofen (ITB) Therapy might be the right choice**, and she completed the trial screening to see how her body would respond.

The trial results were encouraging, reducing Mary's spasticity significantly, and she completed the ITB implant procedure. Mary described that prior to ITB Therapy, she felt handicapped—any pressure on the right side would trigger her spasticity. Since having the pump, however, she no longer uses a wheelchair and has been able to ambulate with a cane. She's also able to take walks in the park with her family and travel, **feeling that she has truly gotten her life back.**

TALK TO YOUR DOCTOR ABOUT MANAGEMENT OPTIONS FOR SPASTICITY RELATED TO A STROKE.

*Patient name and image has been changed to protect their privacy.

SOURCES

1. <http://www.spasticityalliance.org/about-spasticity/stroke-and-spasticity/>, Last accessed: February 7, 2018
2. Bhimani, Rozina and Anderson, Lisa. "Clinical Understanding of Spasticity: Implications for Practice." Rehabilitation Research and Practice Volume 2014, Article ID 279175, 10 pages. <http://dx.doi.org/10.1155/2014/279175>
3. <https://emedicine.medscape.com/article/2207448-overview#a6>
4. Doctor Questionnaire: Stroke Patient Survey. New Jersey. (2019).
5. McGuire JR. Epidemiology of spasticity in the adult and child. In: Brashear A, et al., eds. Spasticity: Diagnosis and Management. New York: Demos Medical Publishing; 2011:5-15.



Piramal, Piramal Critical Care and the Piramal logo are trademarks of a Piramal company.

© 2017 Piramal Critical Care, Inc.
March 2019 MKT-PIR-0059